


GeoForschungsZentrum Potsdam

Geomagnetic Results Wingst

2001

Yearbook No. 47



Potsdam 2004



The Wingst Observatory staff in front of the Office Building. In the background: the Large Laboratory which contains a Fanselau coil system



Wingst Observatory: Fanselau coil system constructed in 1943

Cover: Compass after Pierre de Maricourt, 1269 (SCHÜCK, 1911)

Geomagnetic Results Wingst 2001 – Yearbook No 47

Günter Schulz

Contents

1	Introduction	4
2	General remarks	6
2.1	Recording systems	6
2.2	Levels, standards and constants	7
2.3	Special measurements	8
3	Absolute measurements	8
3.1	Declination and inclination	8
3.2	Horizontal and total intensity, vertical component	9
4	Digital recording system	9
4.1	Base line values	10
4.2	Scale values, temperature, coefficients and cross talk	10
5	Data processing	11
6	Indices	11
7	File set on the CDrom	12
8	References	14
<i>Appendix 1: Figures</i>		
Figure 1	Base line values 2001	15
Figure 2	Daily mean values 2001	16
Figure 3	Epoch values Wingst	17
Figure 4	Files on the CDrom	18
<i>Appendix 2: Tables</i>		
Table 1	Base line values 2001	19
Table 2	Monthly mean values 2001	20
Table 3	Epoch values	21
Table 4	Statistics of indices 2001	22
Tables 5	Hourly mean values 2001	CDrom
Table 6	Indices 2001	CDrom

1 Introduction

This report (yearbook No 47) contains the results of Erdmagnetisches Observatorium Wingst (WNG) for 2001.¹

The enclosed CDrom contains recorded minute values as well as derived (hourly, daily, monthly) mean values and indices. It also provides recalculated epoch values from 1939.5 on and those of Marineobservatorium Wilhelmshaven (WLH) before then. Revised sets of monthly and daily mean values (since 1943) and K values (since 1944) are also included.

Using the visualisation software year.exe, the one-minute, hourly and daily values of the year under review can be displayed as graphs in the same manner as in the years before. High resolution magnetograms for each day are stored as post script files on the CDrom.

In the year under review, Wingst Observatory additionally published on a monthly basis:

- a) Reports on geomagnetic indices and special geomagnetic events
- b) Reports on preliminary daily and monthly means

Geomagnetic data have been provided on a regular basis to the following institutions:

- a) International Space Environment Service (ISES): Geomagnetic indices and geomagnetic events (daily)
- b) International Service of Geomagnetic Indices (ISGI): Geomagnetic indices and special geomagnetic events (monthly and annually)
- c) World Data Centers for Geomagnetism: geomagnetic indices and one-minute values (annually)
- d) INTERMAGNET (Global near-real-time magnetic observatory network): One-minute values (reported data via METEOSAT and Email, hourly; adjusted data via Email, on weekdays); Geomagnetic indices and one-minute values (CDrom, annually)

¹ Reports up to 1999 were published by Bundesamt für Seeschifffahrt und Hydrographie The last one (SCHULZ, 2004) contains a complete digital set of all data that have been published since the establishment of Wingst Observatory in 1938.

Indices and information about special events were made available through a telephone service on weekdays.

Phone: +49 4778 812152

The preliminary variations and indices can be found on the Internet on a real time basis (10 min updates) in graphical form:

http://www.gfz-potsdam.de/pb2/pb23/GeoMag/Other/BothObs_e.html

or (update every hour):

<http://www.bsh.de/en/Marine%20data/Observations/Geomagnetism/obs.jsp>

and in numerical form (update every 10 minutes, pass word required):

<ftp://wng@ftp.bsh.de/outgoing/boulder>

Definitive (compressed) data from 1939 onwards (minute values since 1981) can be found at:

<ftp://ftp.bsh.de/outgoing/wng>

The following list shows some additional selected links providing Wingst data:

Intermagnet (variations):

<http://www.intermagne.bgs.ac.uk/cgi-bin/imagform>

RWC Brussels (indices):

<http://sidc.oma.be/products/wng/index.php3>

WDC Kyoto (pulsations):

<http://swdcft49.kugi.kyoto-u.ac.jp/film/index.html>

WDC Copenhagen (variations):

<http://web.dmi.dk/fsweb/projects/wdcc1/obs.html>

Address for data requests, data exchange and information:

Erdmagnetisches Observatorium
Am Olymp 13
D-21789 Wingst

Phone: +49 4778 812110

Fax: +49 4778 812150

E-mail: guenter.schulz@bsh.de

Collaborators: W.D. Grube and A. Glodek.

2 General Remarks

Wingst Geomagnetic Observatory was established in 1938 as a successor to Wilhelmshaven. Since then, the station has been operated without interruption. The observatory's development is described by VOPPEL, 1988, and SCHULZ, 2001 (see also yearbook No 46, 2000, appendix 3). The development of the modern recording devices is given by SCHULZ, 1998. For the instrumentation since 1938, see also instr.txt on the Cdrom.

The observatory is located in the Lower Elbe area on top of a terminal moraine of the Saale glacial period (elevation 50 m). Its co-ordinates are:

	Latitude	Longitude
Geographic	53° 44.6'N	09° 04.4'E
Geomagnetic	54.2°	95.3°

Geomagnetic co-ordinates refer to DGRF (Definitive Geomagnetic Reference Field) 1980.

The following abbreviations are used throughout this report:

- X North component
- Y East component
- Z Vertical component (downward positive)
- H Horizontal intensity
- D Declination (eastward positive)
- I Inclination (downward positive)
- F Total intensity
- U North-west component
- V North-east component

Times are related to UTC (Co-ordinated Universal Time).

2.1 Recording systems

The results of this edition were derived from the following recording systems:

a) Digital system for variations:

- Suspended fluxgate magnetometer (FM) of type FGE(DMI) (U, V, Z): One-minute and hourly means as well as indices of activity
- Proton precession magnetometer (PPM) of type V75(VARIAN) (F): One minute spot values for quality check only

- b) Photographic system for variations of type SCHULZE/LA COUR (D , H and Z ; 20 mm h^{-1}): Geomagnetic events (ssc, sfe, bay) and substitute hourly means
- c) Photographic system for pulsations of type KIM762(KARMANN) (amplitude and phase characteristics see yearbook No 30, 1984): Geomagnetic events (pc, pi)

2.2 Levels, standards and constants

The results of this edition refer to the International Magnetic Standard (IMS). The results of the yearbooks up to and including 1980 referred to the Observatory Standard (OBS), which was represented by the classic type base line instruments bound to their original locations and surroundings.

H , Z , and F are referred to the proton vector magnetometer (PVM) of type ASKANIA/V4931(VARIAN) on pier NW (section 3.2), D to the fluxgate theodolite (DI-flux) of type 010B(ZEISS)/MAG01H(BARTINGTON) on pier NE (section 3.1) of the absolute house. Both instruments are assumed to represent IMS.

The following equations apply to D (see yearbook No 37, 1991), H and Z (see yearbook No 38, 1992):

$$\begin{aligned} D_{OBS} &= D_{IMS} \\ H_{OBS} &= H_{IMS} + 6.7 \text{ nT} \\ Z_{OBS} &= Z_{IMS} + 11.1 \text{ nT}. \end{aligned}$$

The differences for the derived elements depend on the components, i.e. for 2001:

$$\begin{aligned} F_{OBS} &= F_{IMS} + 12.8 \text{ nT} \\ I_{OBS} &= I_{IMS} - 0.15' \\ X_{OBS} &= X_{IMS} + 6.7 \text{ nT} \\ Y_{OBS} &= Y_{IMS} \end{aligned}$$

The following physical standards are available at Wingst. They guarantee the quality of data:

SCHWILLE (frequency, DCF77, 10^{-8})
 PATEK PHILIPPE and HOPF (UTC, DCF77)
 CROPICO VS10 (Voltage, $5 \cdot 10^{-6}$)
 GUILDLINE 100 Ohm (resistance, $5 \cdot 10^{-6}$)
 Helmholtz coil of high precision (magnetic field strength, 10^{-4})

For the determination of the magnetic induction, the IAGA-recommended gyromagnetic constant (RASMUSSEN, 1991) was used:

$$2\pi\tau^{-1} = 23.487203 \text{ nT s}$$

The azimuth marks were last checked by the German Geodetic Survey in 1995. Their values, related to the NE pier (R: 3504926.873, H: 5956702.028), and their deviations in the year under review are:

Azimuth mark	Azimuth	Deviation against
N	3811° .36'	N
NE	13° 23.19'	(-0.17 ±0.04)'
W	308° 42.94'	(0.01 ±0.04)'

2.3 Special measurements

In the year under review, no comparative measurements were carried out.

3 Absolute measurements

The absolute measurements were reduced according to the variations of the digital system (section 4).

3.1 Declination and Inclination

Absolute measurements of D were made with the DI -flux on an approximately monthly basis. Also the determination of I was included in the measurement routine. Each measurement is based on a set of four positions. I was corrected by the pier difference of $-0.2'$ in the sense of NW minus NE. The differences $E=I-\arctg(Z/H)$ are shown in Table 1.

Additionally, relative measurements of D were carried out with the PVM according to the addition field method (Serson) on a weekly basis. The mean difference in the sense of PVM minus DI -flux of all pairs of measurements carried out on the same day was used as an instrument constant. Its value e is as follows:

$$e = -23.92' \text{ (11 measurements).}$$

3.2 Horizontal intensity, vertical component and total intensity

Absolute measurements of H and Z were carried out with the PVM according to the compensation field method (Nelson) after each relative determination of D .

The magnetic induction vector is over-determined due to the measurement of three elements within the meridian plane. The difference $c = F - (H^2 + Z^2)^{1/2}$ represents the measurements' inherent accuracy. The annual mean of the error C amounted to:

$$+0.2 \text{ nT} \pm 0.4 \text{ nT value (52 measurements).}$$

C is shown in Table 1.

As a rule, the PPM of type V75 was used. This instrument shows a long-term drift of some 0.1 nT depending on the components (SCHULZ AND CARSTENS, 1979). Therefore, comparative measurements using the PPM of type V4931, which represents IMS (see section 2.2), were carried out on a monthly basis. All base line values as well as the recorded minute spot values of F (section 4) are referred to this instrument.

4 Digital recording system

Minute mean values of the orthogonal components U , V , and Z as well as spot values of F were acquired by the primary digital system (V75 and FGE (No 125), section 2.1). The PPM is not only part of the recording system but also serves as an indicator of the PVM (section 3).

Owing to over-determination, outliers, jumps and short-term base line instabilities between the dates of absolute measurements of all three components could be detected (section 4.1) and, under certain conditions, automatically eliminated. The following equation applies to Wingst:

$$dF = 0.26 dU + 0.26 dV + 0.93 dZ.$$

Additionally, a fourth fluxgate was operated, which had been aligned in such a way that its W orientation satisfies the following equation:

$$dW = 0.578(dU + dV + dZ).$$

In this way, jumps and outliers of the secondary system could be monitored independently.

A second suspended FM of type FGE (No 126), an FM of type EDA FM100B and a PPM of type PPM105(EDA) were operated as stand-by devices in case of failure of the primary system.

4.1 Base line values

Table 1 shows the base line values of the FGE125 referred to IMS. Fig 1 shows the results in graphical form. Absolute measurements of D and I (DI-flux) are marked by circles, those of H and Z (PVM) as well as relative measurements of D by dots. I (derived from H and Z) is also displayed (dots).

To obtain base line values, the dependence of the measured elements D , H , I and F on the recorded components U , V , and Z within the range of variations was developed up to terms of second order (see yearbook No 46, 2000, appendix 3). Minute mean values of the magnetometer and the baseline instruments were processed, which had been synchronized within ± 5 s.

For 2001, the base line values of the primary components refer to the following equivalent voltages E of the fluxgate compensation fields:

Component	E in mV (nominal)
U	12861
V	12613
Z	45463

4.2 Scale values, temperature coefficients and cross talk

Scale values and cross talk were traced back to the respective parameters of the old FM100C(EDA) system by employing stochastic methods, making use of strong variations during a substorm on April 7, 1995 (SCHULZ, 1998). The following values apply to the primary components (FGE125):

	Scale Values in nT/mV 1.000+	Cross Talk against FM100C in 10^{-3}	
U	$+10^{-3}(1.4 \pm 0.6)$	$V: +0.2 \pm 1.0$	$Z: +0.9 \pm 0.6$
V	$-10^{-3}(1.5 \pm 0.8)$	$U: -0.7 \pm 0.6$	$Z: -0.5 \pm 0.4$
Z	$+10^{-3}(0.8 \pm 0.6)$	$U: -0.6 \pm 0.4$	$V: -1.2 \pm 0.8$

Considering the respective values of the FM100C (see yearbook No 41, 1995), the absolute misalignments and errors of the scale values of the FGE125 fluxgates probably do not exceed the order of magnitude of 10^{-3} .

Temperature coefficients were neglected because the FGE double system had been installed in the old variometer room (SCHULZ 2001) with almost perfect temperature control (contact thermometers, $\pm 0.03^\circ\text{C}$).

5 Data processing

The base line values (Tables 1) were smoothed by Bathspline approximation in steps of 0.01' for D or 0.1 nT for H and Z , respectively (SCHOTT, 1992).

Hourly mean values were formed using 60 minute mean values of U , V , and Z (taken at minutes 00 to 59 UTC and centred at second 30) as well as 60 F spot values (taken at second 05).

The international quiet (Q) and disturbed (D) days were taken from the Niemegek listings of ISGI. A denotes normal days. In the case of averaging, A means that all days of the month or the year, respectively, have been included.

The data were processed by a computer double system of type HP9000 330/360. Each workstation is connected to a data acquisition unit of type HP3852 and to the Internet. All necessary calculations including those for the yearbook were carried out by the workstation of type HP9000 360.

6 Indices

The indices presented in this edition (File wng01.k and Table 4) indicate the local disturbances of the geomagnetic field resulting from particle radiation. Their meaning in detail:

K : geomagnetic three-hourly index, quasi-logarithmic measure of the maximum disturbance in steps of 0 to 9; lower limit for $K = 9$: 500 nT

sum : Sum of the eight three-hourly indices of a day

Ak : Mean value of the equivalent amplitudes derived from the eight three-hourly indices. The mean value of the daily disturbance of the geomagnetic field is $2 Ak$ nT

Ck : daily character figure derived from Ak and scaled from 0.0 to 2.5.

C : estimated daily character figure; scale: 0, 1, 2

The indices were derived using the IAGA-recommended FMI-routine (Häkkinen, 1992).

7 Files on the CDrom

<i>\wingst\</i>	<i>Wingst root directory, containing the following subdirectories and files</i>
tree_01.txt:	File structure
info.txt:	Information on the operating system
<i>yearb01\:</i>	<i>Directory containing this report (yearbook No 47), tables 5 and 6, magnetograms and a reprint</i>
yearb01\yearb01.pdf:	This report
yearb01\tabs5_01\:	Directory containing tables wngYYmmm.e of hourly and daily mean values for the month mmm of the element e data (<i>D</i> in 0.1', <i>H</i> and <i>Z</i> in nT).
yearb01\tab6_01.txt:	Table of indices
yearb01\mags01\dhz2001mmdd.ps:	Magnetograms (post script) of the day dd in the month mm
yearb01\instr.txt:	Instruments used since 1938
<i>progs\:</i>	<i>Directory containing software</i>
progs\year.exe:	Visualisation programme for hourly and daily mean values as well as one-minute values located in data01
progs\readme.txt:	Notes concerning operation of the programme year.exe and the meaning of the parameters in year.ini
progs\setup.bat:	Installs the programme year under the local directory c:\year and starts the visualisation software
<i>data01\:</i>	<i>Directory containing the following data</i>
data01\wlh+wng.yr:	Updated epoch values WLH and WNG (<i>D</i> and <i>I</i> in 0.1'; <i>F</i> , <i>H</i> , <i>X</i> , <i>Y</i> , and <i>Z</i> in nT)
data01\wng.mon:	Updated monthly mean values WNG (since 1943; <i>D</i> and <i>I</i> in 0.1'; <i>F</i> , <i>H</i> , <i>X</i> , <i>Y</i> , and <i>Z</i> in nT)
data01\wng.day:	Updated daily mean values WNG (since 1944; <i>D</i> and <i>I</i> in 0.1'; <i>F</i> , <i>H</i> , <i>X</i> , <i>Y</i> , and <i>Z</i> in nT)

data01\wng.k: Updated activity figures K , A_k , C_k , and C as well as monthly and annual mean values of A_k , C_k , and C (since 1944)

yearb.exe input files:

data01\hour01\wng01mmm.wdc: Hourly mean values of the month mmm in the format WDC (ICSU, 1989); yearb.exe input files

data01\min01mm\wng01mmm.0nn: One-minute values of the days nn for the month mm or mmm , respectively, in the format WDC (ICSU, 1989); yearb.exe input files

data01\iaga01\:
Directory containing the following data in the IAGA2000 format (IYEMORI et al., 2002). See also: <http://www.ngdc.noaa.gov/IAGA/wg2>

data01\iaga01\YR.WNG: Epoch values WNG starting 1939 (from 1981 on: D and I in 0.01'; X , Y , Z , H and F in 0.1 nT; before then: 0.1' or 1 nT, respectively)

data01\iaga01\2001MT.WNG: Monthly means (D and I in 0.01'; X , Y , Z , H and F in 0.1 nT)

data01\iaga01\2001DY.WNG: Daily means (D and I in 0.01'; X , Y , Z , H and F in 0.1 nT)

data01\iaga01\2001mmHR.WNG: Hourly means (F , X , Y and Z in 0.1 nT) of the month mm

data01\iaga01\2001mmMN.WNG: Minute means (F , X , Y and Z in 0.1 nT) of the month mm

8 References

IYEMORI, T., TAKEDA, M., KAMEI, M. AND M. NOSE, 2002: World Data Center for Geomagnetism, Kyoto, Data catalogue No 26

ICSU, 1989: Guide to the World Data Center System, Part 3, Geomagnetism; ICSU Panel of WDCs, Boulder

HÄKKINEN, L., 1992: Routine for computing daily K-indices by the FMI-method, Fortran source. Recommended by the IAGA WG V-5, 1993, IAGA News, No 32, 27

RASMUSSEN, O., 1991: The proton gyromagnetic ratio. IAGA News, No 30, 78

SCHOTT, J. J., 1992: Personal communication

Schück, A., 1911: Der Kompass, Tafel 1. Hamburg

SCHULZ, G. and U. CARSTENS, 1979: A period measuring proton magnetometer with a direct readout. Dt. hydrogr. Z. 32, 119-125

SCHULZ, G., 1998: Long-term experience with variometer systems of different generations at Wingst observatory. GFZ Scient.Tech.Report 98/21

SCHULZ, G., 2001: From Deutsche Seewarte Hamburg to GeoForschungsZentrum Potsdam – Wingst Geomagnetic Observatory during six decades. Contr. to Geophysics and Geodesy 31, 17-24

SCHULZ, G., 2004: Geomagnetic Results Wingst, 1996 1997 1998 1999 including the complete Wingst data set since 1939 on CDrom. Berichte des BSH, No 33

VOPPEL, D., 1988: Some remarks on the history of Wingst Geomagnetic Observatory during the first 50 years. Dt. hydrogr. Z. 41, 109-117

Appendix 1 and 2: Figures and Tables




Fig. 1

Wingst 2001 Base line values of the fluxgate system FGE125, IMS




Fig. 2

Wingst 2001 Daily mean values D , H and Z




Fig. 3

Wingst Epoch values I, D, F, H and Z

```

wingst\
  tree_01.txt
  info.txt
  yearb01\
    yearb01.pdf
    tabs5_01\
      wng01jan.d
      wng01jan.h
      wng01jan.z
      .
      wng01dec.d
      wng01dec.h
      wng01dec.z
    tab6_01.txt
    mags01\dhz20010101.ps
    .
    mags01\dhz20011231.ps
    instr.txt
  progs\
    readme.txt
    year.exe
    year.ini
    setup.bat
    att.bgi
    cga.bgi
    egavga.bgi
    herc.bgi
    vesal6.bgi
  data01\
    wlh+wng.yr
    wng.mon
    wng.day
    wng.k
    hour01\
      wng01jan.wdc
      .
      wng01dec.wdc
    min0101\
      wng01jan.001
      .
      wng01jan.031
      .
    min0112\
      wng01dec.001
      .
      wng01dec.031
    iaga01\
      YR.WNG
      2001MT.WNG
      2001DY.WNG
      200101MT.WNG
      .
      200112MT.WNG
      200101HR.WNG
      .
      200112HR.WNG

```

Fig. 4

Structure of the file set on CDrom

Wingst 2001

Base-line measurements, system FGE125, IMS

Month	day	Do(abs)	Do(rel)	Io	Fo nT	Ho nT	Zo nT	C	E
Jan.	3		-0°17.46'		48960.4	18030.1	45519.3	+0.3	
	10		-0 17.63		48961.1	18029.8	45519.6	+0.8	
	17		-0 17.39		48961.0	18030.1	45519.5	+0.7	
	24		-0 17.28		48961.0	18031.1	45519.9	+0.0	
	31	-0°17.45'	-0 17.48	+68°23.71'	48960.9	18031.0	45519.9	-0.1	+0.26'
Feb.	7		-0 17.10		48960.8	18028.9	45520.0	+0.5	
	14		-0 17.29		48961.2	18030.8	45519.7	+0.5	
	21		-0 17.57		48960.7	18030.3	45519.6	+0.2	
	28	-0 17.39	-0 17.57	+68 23.68	48961.1	18030.6	45519.8	+0.4	+0.20
March	7		-0 17.33		48961.3	18030.8	45519.7	+0.6	
	15		-0 17.44		48961.2	18031.0	45519.9	+0.2	
	21		-0 17.39		48961.3	18031.2	45519.4	+0.7	
	29		-0 17.54		48961.2	18031.0	45519.9	+0.2	
April	4	-0 17.41	-0 17.24	+68 23.62	48960.6	18031.6	45519.4	-0.1	+0.22
	11		-0 17.28		48961.2	18031.4	45520.3	-0.3	
	18		-0 17.38		48960.6	18031.2	45519.7	-0.3	
	24	-0 17.25	-0 17.33	+68 23.59	48960.9	18030.8	45519.3	+0.5	+0.14
May	2		-0 17.40		48960.4	18030.9	45519.4	-0.1	
	9		-0 17.43		48960.6	18029.9	45520.1	-0.2	
	16		-0 17.34		48960.5	18030.8	45519.6	-0.1	
	23	-0 17.35	-0 17.35	+68 23.67	48960.6	18030.5	45519.4	+0.3	+0.20
	30		-0 17.51		48960.4	18030.0	45519.2	+0.4	
June	6		-0 17.40		48960.7	18030.2	45519.8	+0.1	
	13		-0 17.48		48960.6	18031.2	45519.4	+0.0	
	20		-0 17.39		48960.0	18030.0	45519.4	-0.2	
	27	-0 17.18	-0 17.25	+68 23.67	48960.2	18030.5	45519.3	+0.0	+0.20
July	4		-0 17.29		48960.1	18029.7	45519.5	+0.0	
	11		-0 17.27		48960.4	18030.4	45519.3	+0.2	
	18		-0 17.47		48960.4	18031.1	45519.1	+0.1	
	25	-0 17.36	-0 17.05	+68 23.72	48959.9	18030.2	45518.4	+0.6	+0.25
Aug.	1		-0 17.27		48960.7	18030.5	45518.9	+0.8	
	8		-0 17.05		48960.6	18030.9	45518.9	+0.6	
	15		-0 17.19		48960.5	18029.9	45518.5	+1.2	
	22		-0 17.29		48960.8	18030.7	45519.1	+0.7	
	29	-0 17.11	-0 17.32	+68 23.56	48960.0	18029.9	45518.4	+0.8	+0.08
Sep.	6		-0 17.30		48960.5	18031.2	45518.8	+0.5	
	12		-0 17.27		48960.3	18030.7	45518.3	+0.9	
	19		-0 17.19		48960.5	18030.8	45519.0	+0.4	
	26		-0 17.29		48960.3	18030.5	45519.8	-0.4	
Oct.	4	-0 17.04	-0 17.26	+68 23.62	48960.0	18030.0	45519.3	-0.1	+0.12
	10		-0 17.27		48960.0	18030.2	45519.2	+0.0	
	17		-0 17.23		48960.0	18029.2	45519.6	+0.0	
	24		-0 17.25		48959.9	18029.8	45519.3	-0.1	
	31		-0 17.06		48959.7	18029.9	45519.1	-0.1	
Nov.	7	-0 17.20	-0 17.08	+68 23.72	48959.9	18030.2	45519.5	-0.4	+0.23
	14		-0 17.10		48960.4	18030.4	45520.0	-0.5	
	21		-0 17.17		48960.3	18030.0	45519.9	-0.3	
	28		-0 17.22		48960.8	18030.3	45520.6	-0.6	
Dec.	5	-0 17.36	-0 17.21	+68 23.75	48960.9	18030.7	45520.0	-0.1	+0.28
	12		-0 17.24		48960.9	18030.6	45520.1	-0.1	
	19		-0 17.10		48960.7	18030.4	45520.0	-0.2	
	27		-0 17.14		48960.5	18030.6	45519.7	-0.2	

Table 1 Wingst 2001 base line values of the fluxgate system FGE125

Wingst (WNG)
 Geographic Coordinates: 53.743° N 9.073° E

2001

Monthly mean values, IMS

D: disturbed, Q: quiet, A: all days

Month		D	F nT	H nT	I	X nT	Y nT	Z nT
Jan	A	-18.8'	49151	18073	68°25.5'	18073	99	45707
Feb	A	-19.4	49151	18077	68 25.2	18077	102	45706
Mar	A	-20.5	49153	18068	68 25.9	18068	108	45711
Apr	A	-21.9	49162	18064	68 26.5	18064	115	45723
May	A	21.4	49162	18086	68 24.9	18086	113	45714
Jun	A	21.9	49164	18088	68 24.8	18088	115	45716
Jul	A	22.4	49165	18087	68 24.9	18087	118	45717
Aug	A	22.9	49168	18083	68 25.3	18082	121	45722
Sep	A	23.8	49169	18077	68 25.8	18076	125	45725
Oct	A	25.4	49180	18056	68 27.6	18056	133	45746
Nov	A	26.0	49185	18060	68 27.5	18059	136	45750
Dec	A	25.5	49188	18077	68 26.3	18076	134	45746
Mean	A	22.5	49167	18075	68 25.9	18074	118	45724
Jan	Q	18.5	49150	18081	68 24.9	18081	97	45703
Feb	Q	19.1	49151	18081	68 25.0	18081	101	45705
Mar	Q	19.7	49152	18083	68 24.8	18083	104	45704
Apr	Q	21.3	49162	18076	68 25.6	18076	112	45718
May	Q	21.2	49161	18088	68 24.7	18088	112	45713
Jun	Q	22.2	49165	18089	68 24.8	18089	117	45717
Jul	Q	22.1	49165	18089	68 24.8	18089	116	45716
Aug	Q	23.0	49168	18084	68 25.2	18083	121	45722
Sep	Q	22.9	49167	18084	68 25.2	18084	121	45721
Oct	Q	24.7	49179	18072	68 26.4	18071	130	45739
Nov	Q	25.2	49187	18074	68 26.5	18073	133	45746
Dec	Q	25.5	49187	18078	68 26.2	18077	134	45744
Mean	Q	22.1	49166	18082	68 25.3	18081	117	45721
Jan	D	19.4	49152	18062	68 26.5	18062	102	45714
Feb	D	20.1	49150	18071	68 25.8	18070	106	45708
Mar	D	22.8	49155	18033	68 28.6	18033	120	45727
Apr	D	23.6	49159	18046	68 27.9	18045	124	45727
May	D	22.2	49161	18075	68 25.7	18075	117	45718
Jun	D	22.2	49163	18082	68 25.1	18082	116	45716
Jul	D	22.5	49165	18086	68 25.0	18085	119	45718
Aug	D	23.1	49171	18079	68 25.6	18079	121	45726
Sep	D	25.1	49170	18063	68 26.8	18063	132	45732
Oct	D	27.6	49180	18027	68 29.8	18026	145	45757
Nov	D	28.2	49177	18018	68 30.4	18017	148	45757
Dec	D	26.1	49186	18067	68 27.0	18066	137	45748
Mean	D	23.6	49166	18059	68 27.0	18059	124	45729

Table 2 Monthly and annual mean values 2001

Wingst (WNG)

Geographic Coordinates: 53.743°N 9.073°E

Annual mean values (IMS)

Epoch	D	F nT	H nT	I	X nT	Y nT	Z nT
1939.5	-5°59.1'	47476	17630	68°12.1'	17534	-1838	44081
1940.5	-5 50.2	47506	17624	68 13.4	17533	-1792	44116
1941.5	-5 40.8	47550	17617	68 15.2	17530	-1744	44166
1942.5	-5 33.1	47579	17622	68 15.7	17540	-1705	44196
1943.5	-5 24.2	47634	17614	68 18.0	17535	-1659	44259
1944.5	-5 16.2	47652	17616	68 18.3	17541	-1618	44276
1945.5	-5 8.2	47671	17611	68 19.2	17540	-1577	44299
1946.5	-4 59.6	47708	17595	68 21.5	17528	-1532	44346
1947.5	-4 51.7	47726	17596	68 22.0	17532	-1491	44365
1948.5	-4 44.4	47775	17602	68 22.9	17541	-1454	44415
1949.5	-4 36.6	47791	17604	68 23.2	17547	-1415	44431
1950.5	-4 29.1	47814	17617	68 22.9	17562	-1378	44451
1951.5	-4 21.5	47832	17624	68 22.8	17573	-1339	44468
1952.5	-4 14.5	47861	17636	68 22.7	17587	-1304	44494
1953.5	-4 7.6	47882	17653	68 22.0	17607	-1270	44510
1954.5	-4 1.3	47899	17666	68 21.5	17623	-1239	44523
1955.5	-3 55.1	47930	17676	68 21.6	17634	-1208	44552
1956.5	-3 49.3	47964	17676	68 22.6	17636	-1178	44589
1957.5	-3 44.2	47993	17686	68 22.6	17648	-1152	44616
1958.5	-3 39.5	48023	17700	68 22.4	17663	-1129	44643
1959.5	-3 34.6	48062	17714	68 22.4	17679	-1105	44679
1960.5	-3 30.1	48095	17727	68 22.4	17693	-1082	44710
1961.5	-3 25.7	48117	17751	68 21.1	17719	-1061	44723
1962.5	-3 21.3	48136	17773	68 20.0	17742	-1040	44735
1963.5	-3 16.9	48160	17789	68 19.4	17760	-1018	44755
1964.5	-3 13.1	48183	17810	68 18.4	17782	-1000	44771
1965.5	-3 9.6	48201	17829	68 17.5	17802	-983	44783
1966.5	-3 6.3	48226	17842	68 17.3	17815	-966	44805
1967.5	-3 3.4	48254	17855	68 17.1	17829	-952	44830
1968.5	-3 1.0	48286	17874	68 16.5	17849	-941	44857
1969.5	-2 59.2	48320	17899	68 15.5	17874	-932	44883
1970.5	-2 56.9	48359	17924	68 14.7	17900	-922	44915
1971.5	-2 54.5	48397	17953	68 13.6	17930	-911	44944
1972.5	-2 51.0	48434	17977	68 12.9	17954	-894	44975
1973.5	-2 46.6	48473	17999	68 12.2	17978	-872	45008
1974.5	-2 41.4	48513	18018	68 11.9	17998	-846	45043
1975.5	-2 36.0	48549	18043	68 11.0	18024	-818	45073
1976.5	-2 29.3	48583	18062	68 10.5	18045	-784	45101
1977.5	-2 22.4	48612	18078	68 10.1	18062	-748	45126
1978.5	-2 14.1	48646	18081	68 10.9	18066	-705	45161
1979.5	-2 6.3	48668	18089	68 10.9	18076	-664	45181
1980.5	-1 59.0	48682	18096	68 10.7	18085	-626	45194
1981.5	-1 51.4	48704	18091	68 11.7	18082	-586	45220
1982.5	-1 43.9	48724	18084	68 12.8	18076	-546	45244
1983.5	-1 36.9	48738	18087	68 13.0	18080	-510	45257
1984.5	-1 29.9	48752	18083	68 13.7	18077	-473	45274
1985.5	-1 23.5	48768	18080	68 14.4	18075	-439	45292
1986.5	-1 17.0	48787	18071	68 15.5	18067	-404	45316
1987.5	-1 11.1	48804	18069	68 16.2	18065	-374	45336
1988.5	-1 5.0	48829	18056	68 17.9	18053	-341	45368
1989.5	-59.0	48856	18042	68 19.7	18039	-309	45402
1990.5	-53.9	48875	18041	68 20.3	18038	-283	45423
1991.5	-48.5	48895	18032	68 21.5	18031	-255	45448
1992.5	-43.4	48911	18038	68 21.5	18037	-228	45463
1993.5	-37.1	48928	18044	68 21.6	18043	-195	45479
1994.5	-30.0	48952	18045	68 22.2	18044	-158	45505
1995.5	-23.0	48975	18053	68 22.2	18053	-121	45526
1996.5	-15.6	48998	18062	68 22.1	18062	-82	45547
1997.5	-7.6	49028	18063	68 22.9	18063	-40	45579
1998.5	0.5	49062	18059	68 24.2	18059	3	45618
1999.5	8.0	49094	18063	68 24.7	18063	42	45651
2000.5	15.4	49132	18064	68 25.7	18064	81	45690
2001.5	22.5	49167	18075	68 25.9	18074	118	45724

Table 3 Wingst Epoch values from 1939 to 2001

Wingst (WNG)

Geographic Coordinates: 53.743° N 9.073° E

2001

Absolute and relative Frequencies of the Three-hourly Index K

K	UTC	0-3	3-6	6-9	9-12	12-15	15-18	18-21	21-24
0		27	26	20	7	12	23	14	16
1		79	83	103	76	67	63	65	69
2		106	140	150	150	142	105	102	101
3		89	82	63	93	94	89	100	104
4		43	22	21	30	30	54	40	45
5		15	6	5	5	15	19	29	17
6		4	3	1	4	4	8	10	6
7		0	2	0	0	1	2	5	6
8		1	1	2	0	0	2	0	0
9		1	0	0	0	0	0	0	1

Absolute Number of Days during the Year for a given K

K	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
0	23	30	8	6	3	4	2	5	9	13	31	11	145
1	75	72	46	28	59	44	29	36	49	40	71	56	605
2	75	71	88	74	90	88	105	86	84	61	72	102	996
3	49	39	48	69	59	64	81	79	59	77	33	57	714
4	18	9	31	28	23	33	27	31	24	28	14	19	285
5	6	3	14	23	12	6	4	9	9	15	7	3	111
6	2	0	7	5	2	1	0	2	5	9	7	0	40
7	0	0	4	4	0	0	0	0	1	5	2	0	16
8	0	0	2	2	0	0	0	0	0	0	2	0	6
9	0	0	0	1	0	0	0	0	0	0	1	0	2

Absolute Number of Three-hour-intervals for a given K

Table 4 Statistics of indices 2001